

Slant Riemannian maps to Kähler manifolds

Bayram Şahin

*Inonu University, Department of Mathematics, 44280, Malatya-Turkey.
E-mail: bayram.sahin@inonu.edu.tr*

Abstract. We introduce slant Riemannian maps from Riemannian manifolds to almost Hermitian manifolds as a generalization of slant immersions, invariant Riemannian maps and anti-invariant Riemannian maps. We give examples, obtain characterizations and investigate the harmonicity of such maps. We also obtain necessary and sufficient conditions for slant Riemannian maps to be totally geodesic. Moreover we relate the notion of slant Riemannian maps to the notion of pseudo horizontally weakly conformal maps which are useful for proving various complex-analytic properties of stable harmonic maps from complex projective space.

2000 Mathematics Subject Classification: 53C15, 53B20, 53C43.

Keywords: Riemannian map, Hermitian manifold, Slant immersion, invariant Riemannian map, Anti-invariant Riemannian map, Slant Riemannian map.

1. Introduction

Smooth maps between Riemannian manifolds are useful for comparing geometric structures between two manifolds. Isometric immersions (Riemannian submanifolds) are basic such maps between Riemannian manifolds and they are characterized by their Riemannian metrics and Jacobian matrices. More precisely, a smooth map $F : (M_1, g_1) \longrightarrow (M_2, g_2)$ between Riemannian manifolds (M_1, g_1) and (M_2, g_2) is called an isometric immersion if F_* is injective and

$$g_2(F_*X, F_*Y) = g_1(X, Y) \quad (1.1)$$

for X, Y vector fields tangent to M_1 , here F_* denotes the derivative map.

Let $\bar{M}$ be a Kähler manifold with complex structure J and M a Riemannian manifold isometrically immersed in $\bar{M}$. We note that many types of submanifolds can be defined depending on the behaviour of the tangent bundle of the submanifold under the action of the complex structure of the ambient manifold. A submanifold M is called holomorphic (complex) if $J(T_p M) \subset T_p M$, for every $p \in M$, where $T_p M$ denotes the tangent space to M at the point p . M is called totally real if $J(T_p M) \subset T_p M^\perp$ for every $p \in M$, where $T_p M^\perp$ denotes the normal space to M at the point p . As a generalization of holomorphic and totally real submanifolds, slant submanifolds were introduced by Chen in [7]. We recall that the submanifold M is called slant if for all non-zero vector X tangent to M the angle $\theta(X)$ between JX and $T_p M$ is a constant, i.e, it does not depend on the choice of $p \in M$ and $X \in T_p M$.

On the other hand, Riemannian submersions between Riemannian manifolds were studied by O'Neill [19] and Gray [13]. Later such submersions were considered between manifolds with differentiable structures. As an analogue of holomorphic submanifolds, Watson defined almost Hermitian submersions between almost Hermitian manifolds and he showed that the base manifold and each fiber have the same kind of structure as the total space, in most cases [22]. We note that almost Hermitian submersions have been extended to the almost contact manifolds [9], locally conformal Kähler manifolds [17] and quaternion Kähler manifolds [14] (see: [10] for details concerning Riemannian submersions between Riemannian manifolds equipped with additional structures of complex, contact, locally conformal or quaternion Kähler).

In 1992, Fischer introduced Riemannian maps between Riemannian manifolds in [11] as a generalization of the notions of isometric immersions and Riemannian submersions. Let $F : (M_1, g_1) \longrightarrow (M_2, g_2)$ be a smooth map between Riemannian manifolds such that $0 < \text{rank} F < \min\{m, n\}$, where $\dim M_1 = m$ and $\dim M_2 = n$. Then we denote the kernel space of F_* by $\ker F_*$ and consider the orthogonal complementary space $\mathcal{H} = (\ker F_*)^\perp$ to $\ker F_*$. Then the tangent bundle of M_1 has the following decomposition

$$TM_1 = \ker F_* \oplus \mathcal{H}.$$

We denote the range of F_* by $\text{range} F_*$ and consider the orthogonal complementary space $(\text{range} F_*)^\perp$ to $\text{range} F_*$ in the tangent bundle TM_2 of M_2 . Since $\text{rank} F < \min\{m, n\}$, we always have $(\text{range} F_*)^\perp$. Thus the tangent

bundle TM_2 of M_2 has the following decomposition

$$TM_2 = (range F_*) \oplus (range F_*)^\perp.$$

Now, a smooth map $F : (M_1^m, g_1) \longrightarrow (M_2^n, g_2)$ is called a Riemannian map at $p_1 \in M$ if the horizontal restriction $F_{*p_1}^h : (ker F_{*p_1})^\perp \longrightarrow (range F_{*p_1})$ is a linear isometry between the inner product spaces $((ker F_{*p_1})^\perp, g_1(p_1)|_{(ker F_{*p_1})^\perp})$ and $(range F_{*p_1}, g_2(p_2)|_{range F_{*p_1}})$, $p_2 = F(p_1)$. Therefore Fischer stated in [11] that a Riemannian map is a map which is as isometric as it can be. In another words, F_* satisfies the equation (1.1) for X, Y vector fields tangent to $\mathcal{H}$. It follows that isometric immersions and Riemannian submersions are particular Riemannian maps with $ker F_* = \{0\}$ and $(range F_*)^\perp = \{0\}$. It is known that a Riemannian map is a subimmersion [11]. It is also important to note that Riemannian maps satisfy the eikonal equation which is a bridge between geometric optics and physical optics. For Riemannian maps and their applications in spacetime geometry see: [12].

In [20], we introduced invariant and anti-invariant Riemannian maps as a generalization of holomorphic submanifolds and totally real submanifolds of almost Hermitian manifolds. We gave examples of such maps and obtained fundamental properties of invariant and anti-invariant Riemannian maps. As a generalization of invariant and anti-invariant Riemannian maps, semi-invariant Riemannian maps from Riemannian manifolds to almost Hermitian manifolds were defined and the geometry of such maps was studied in [21].

In this paper, we introduce slant Riemannian maps from Riemannian manifolds to almost Hermitian manifolds. We show that slant Riemannian maps include slant immersions (therefore holomorphic immersions and totally real immersions), invariant Riemannian maps and anti-invariant Riemannian maps. We also obtain an example which is not included in immersions, or invariant Riemannian maps or anti-invariant Riemannian maps. We investigate the harmonicity of slant Riemannian maps and obtain necessary and sufficient conditions for such maps to be totally geodesic. We also show that every slant Riemannian map is a pseudo horizontally weakly conformal map, then we obtain necessary and sufficient conditions for slant Riemannian maps to be a pseudo homothetic map. We note that the notion of pseudo horizontally weakly conformal maps were introduced in [5] to study the stability of harmonic maps into irreducible Hermitian symmetric spaces of compact type, later such maps have been studied in [2], [3] and [16].

2. Preliminaries

In this section, we recall basic facts about Riemannian maps and almost Hermitian manifolds. Let (M, g_M) and (N, g_N) be Riemannian manifolds and suppose that $F : M \rightarrow N$ is a smooth map between them. Then the differential F_* of F can be viewed as a section of the bundle $Hom(TM, F^{-1}TN) \rightarrow M$, where $F^{-1}TN$ is the pullback bundle which has fibres $(F^{-1}TN)_p = T_{F(p)}N, p \in M$. $Hom(TM, F^{-1}TN)$ has a connection ∇ induced from the Levi-Civita connection ∇^M and the pullback connection. Then the second fundamental form of F is given by

$$(\nabla F_*)(X, Y) = \nabla_X^F F_*(Y) - F_*(\nabla_X^M Y) \quad (2.1)$$

for $X, Y \in \Gamma(TM)$, where ∇^F is the pullback connection. It is known that the second fundamental form is symmetric. Let $F : (M, g_M) \rightarrow (N, g_N)$ be a smooth map between Riemannian manifolds and assume that M is compact, then its energy is

$$E(F) = \int_M e(F) v_g = \frac{1}{2} \int_M |dF|^2 v_g.$$

The critical points of E are called harmonic maps. Standard arguments yield the associated Euler-Lagrange equation, the vanishing of the tension field: $\tau(F) = \text{trace}(\nabla F_*)$ [4].

From now on, for simplicity, we denote by ∇^2 both the Levi-Civita connection of (M_2, g_2) and its pullback along F . Then according to [18], for any vector field X on M_1 and any section V of $(\text{range } F_*)^\perp$, where $(\text{range } F_*)^\perp$ is the subbundle of $F^{-1}(TM_2)$ with fiber $(F_*(T_p M))^\perp$ -orthogonal complement of $F_*(T_p M)$ for g_2 over p , we have $\nabla_X^{F^\perp} V$ which is the orthogonal projection of $\nabla_X^2 V$ on $(F_*(TM))^\perp$. In [18], the author also showed that $\nabla^{F^\perp}$ is a linear connection on $(F_*(TM))^\perp$ such that $\nabla^{F^\perp} g_2 = 0$. We now define $\mathcal{S}_V$ as

$$\nabla_{F_* X}^2 V = -\mathcal{S}_V F_* X + \nabla_X^{F^\perp} V, \quad (2.2)$$

where $\mathcal{S}_V F_* X$ is the tangential component (a vector field along F) of $\nabla_{F_* X}^2 V$. It is easy to see that $\mathcal{S}_V F_* X$ is bilinear in V and $F_* X$ and $\mathcal{S}_V F_* X$ at p depends only on V_p and $F_{*p} X_p$. By direct computations, we obtain

$$g_2(\mathcal{S}_V F_* X, F_* Y) = g_2(V, (\nabla F_*)(X, Y)), \quad (2.3)$$

for $X, Y \in \Gamma((\ker F_*)^\perp)$ and $V \in \Gamma((\text{range } F_*)^\perp)$. Since (∇F_*) is symmetric, it follows that $\mathcal{S}_V$ is a symmetric linear transformation of $\text{range } F_*$.

A $2k$ -dimensional Riemannian manifold $(\bar{M}, \bar{g}, \bar{J})$ is called an almost Hermitian manifold if there exists a tensor field $\bar{J}$ of type $(1,1)$ on $\bar{M}$ such that $\bar{J}^2 = -I$ and

$$\bar{g}(X, Y) = \bar{g}(\bar{J}X, \bar{J}Y), \forall X, Y \in \Gamma(T\bar{M}), \quad (2.4)$$

where I denotes the identity transformation of $T_p\bar{M}$. Consider an almost Hermitian manifold $(\bar{M}, \bar{J}, \bar{g})$ and denote by $\bar{\nabla}$ the Levi-Civita connection on $\bar{M}$ with respect to $\bar{g}$, then $\bar{M}$ is called a Kähler manifold [23] if $\bar{J}$ is parallel with respect to $\bar{\nabla}$, i.e.,

$$(\bar{\nabla}_X \bar{J})Y = 0 \quad (2.5)$$

for $X, Y \in \Gamma(T\bar{M})$.

3. Slant Riemannian maps

In this section, we define slant Riemannian maps from a Riemannian manifold to an almost Hermitian manifold. We give examples, obtain characterizations and investigate the harmonicity of a slant Riemannian map. Then we give necessary and sufficient conditions for a slant Riemannian map to be totally geodesic.

Definition 3.1. *Let F be a Riemannian map from a Riemannian manifold (M_1, g_1) to an almost Hermitian manifold (M_2, g_2, J) . If for any non-zero vector $X \in \Gamma(\ker F_*)^\perp$, the angle $\theta(X)$ between $JF_*(X)$ and the space $\text{range } F_*$ is a constant, i.e. it is independent of the choice of the point $p \in M_1$ and choice of the tangent vector $F_*(X)$ in $\text{range } F_*$, then we say that F is a slant Riemannian map. In this case, the angle θ is called the slant angle of the slant Riemannian map.*

Since F is a subimmersion, it follows that the rank of F is constant on M_1 , then the rank theorem for functions implies that $\ker F_*$ is an integrable subbundle of TM_1 , ([1], page:205).

We first give some examples of slant Riemannian maps.

Example 1. Every θ -slant immersion from a Riemannian manifold to an almost Hermitian manifold is a slant Riemannian map with $\ker F_* = \{0\}$.

Example 2. Every invariant Riemannian map from a Riemannian manifold to an almost Hermitian manifold is a slant Riemannian map with $\theta = 0$.

Example 3. Every anti-invariant Riemannian map from a Riemannian manifold to an almost Hermitian manifold is a slant Riemannian map with $\theta = \frac{\pi}{2}$.

A slant Riemannian map is said to be proper if it is not an immersion and $\theta \neq 0, \frac{\pi}{2}$. Here is an example of proper slant Riemannian maps.

Example 4. Consider the following Riemannian map given by

$$F : \begin{array}{ccc} R^4 & \longrightarrow & R^4 \\ (x_1, x_2, x_3, x_4) & & (x_1, \frac{x_2+x_3}{\sqrt{3}}, \frac{x_2+x_3}{\sqrt{6}}, 0). \end{array}$$

Then $\text{rank} F = 2$ and for any $0 < \alpha < \frac{\pi}{2}$, F is a slant Riemannian map with slant angle $\cos^{-1}(\sqrt{\frac{2}{3}})$.

We also have the following result which is based on the fact that the composition of a Riemannian submersion F_1 from a Riemannian manifold (M_1, g_1) onto a Riemannian manifold (M_2, g_2) and an isometric immersion F_2 from (M_2, g_2) to a Riemannian manifold (M_3, g_3) is a Riemannian map.

Proposition 3.1. *Let F_1 be a Riemannian submersion from a Riemannian manifold (M_1, g_1) onto a Riemannian manifold (M_2, g_2, J) and F_2 a slant immersion from (M_2, g_2, J) to an almost Hermitian manifold (M_3, g_3, J) . Then $F_2 \circ F_1$ is a slant Riemannian map.*

Let F be a Riemannian map from a Riemannian manifold (M_1, g_1) to an almost Hermitian manifold (M_2, g_2, J) . Then for $F_*(X) \in \Gamma(\text{range} F_*)$, $X \in \Gamma((\ker F_*)^\perp)$, we write

$$JF_*(X) = \phi F_*(X) + \omega F_*(X), \quad (3.1)$$

where $\phi F_*(X) \in \Gamma(\text{range} F_*)$ and $\omega F_*(X) \in \Gamma((\text{range} F_*)^\perp)$. Also for $V \in \Gamma((\text{range} F_*)^\perp)$, we have

$$JV = \mathcal{B}V + \mathcal{C}V, \quad (3.2)$$

where $\mathcal{B}V \in \Gamma(\text{range} F_*)$ and $\mathcal{C}V \in \Gamma((\text{range} F_*)^\perp)$. We now recall from [20] that the second fundamental form $(\nabla F_*)(X, Y)$, $\forall X, Y \in \Gamma((\ker F_*)^\perp)$, of a Riemannian map has no components in $\text{range} F_*$, i.e.

$$(\nabla F_*)(X, Y) \in \Gamma((\text{range} F_*)^\perp). \quad (3.3)$$

Let F be a Riemannian map from a Riemannian manifold (M_1, g_1) to a Kähler manifold (M_2, g_2, J) , then from (3.1), (3.2), (3.3) and (2.1) we obtain

$$(\tilde{\nabla}_X \omega)F_*(Y) = \mathcal{C}(\nabla F_*)(X, Y) - (\nabla F_*)(X, Y') \quad (3.4)$$

and

$$F_*(\nabla_X^1 Y') - \phi F_*(\nabla_X^1 Y) = \mathcal{B}(\nabla F_*)(X, Y) + \mathcal{S}_{\omega F_*(Y)} F_*(X) \quad (3.5)$$

for $X, Y \in \Gamma((\ker F_*)^\perp)$, where $(\tilde{\nabla}_X \omega)F_*(Y) = \nabla_X^{F^\perp} F_*(Y) - \omega F_*(\nabla_X^1 Y)$, ∇^1 is the Levi-Civita connection of M_1 and $\phi F_*(Y) = F_*(Y')$, $Y' \in \Gamma((\ker F_*)^\perp)$.

Let F be a slant Riemannian map from a Riemannian manifold (M_1, g_1) to an almost Hermitian manifold (M_2, g_2, J) , then we say that ω is parallel if $(\tilde{\nabla}_X \omega)F_*(Y) = 0$. We also say that ϕ is parallel if $F_*(\nabla_X^1 Y') - \phi F_*(\nabla_X^1 Y) = 0$ for $X, Y \in \Gamma((\ker F_*)^\perp)$ such that $\phi F_*(Y) = F_*(Y')$, $Y' \in \Gamma((\ker F_*)^\perp)$.

The proof of the following result is exactly same with slant immersions (see [8] or [6] for Sasakian case), therefore we omit its proof.

Theorem 3.1. *Let F be a Riemannian map from a Riemannian manifold (M_1, g_1) to an almost Hermitian manifold (M_2, g_2, J) . Then F is a slant Riemannian map if and only if there exists a constant $\lambda \in [-1, 0]$ such that*

$$\phi^2 F_*(X) = \lambda F_*(X)$$

for $X \in \Gamma((\ker F_*)^\perp)$. If F is a slant Riemannian map, then $\lambda = -\cos^2 \theta$.

By using above theorem, it is easy to see that

$$g_2(\phi F_*(X), \phi F_*(Y)) = \cos^2 \theta g_1(X, Y) \quad (3.6)$$

$$g_2(\omega F_*(X), \omega F_*(Y)) = \sin^2 \theta g_1(X, Y) \quad (3.7)$$

for any $X, Y \in \Gamma((\text{range } F_*)^\perp)$.

We now recall the notion of adjoint map which will be useful for the results obtained in this section (for more details see [12]). Let $F : (M_1, g_1) \rightarrow (M_2, g_2)$ be a Riemannian map between Riemannian manifolds (M_1, g_1) and (M_2, g_2) . Then the adjoint map ${}^*F_*$ of F_* is characterized by $g_1(x, {}^*F_{*p_1} y) = g_2(F_{*p_1} x, y)$ for $x \in T_{p_1} M_1$, $y \in T_{F(p_1)} M_2$ and $p_1 \in M_1$. Considering F_*^h at each $p_1 \in M_1$ as a linear transformation

$$F_{*p_1}^h : ((\ker F_*)^\perp(p_1), g_1|_{(\ker F_*)^\perp(p_1)}) \rightarrow (\text{range } F_*(p_2), g_2|_{\text{range } F_*(p_2)}),$$

we will denote the adjoint of F_*^h by ${}^*F_{*p_1}^h$. Let ${}^*F_{*p_1}$ be the adjoint of $F_{*p_1} : (T_{p_1}M_1, g_{1p_1}) \longrightarrow (T_{p_2}M_2, g_{2p_2})$. Then the linear transformation

$$({}^*F_{*p_1})^h : \text{range} F_*(p_2) \longrightarrow (\ker F_*)^\perp(p_1)$$

defined by $({}^*F_{*p_1})^h y = {}^*F_{*p_1} y$, where $y \in \Gamma(\text{range} F_{*p_1})$, $p_2 = F(p_1)$, is an isomorphism and $(F_{*p_1}^h)^{-1} = ({}^*F_{*p_1})^h = ({}^*F_{*p_1}^h)$.

Let $\{e_1, \dots, e_n\}$ be an orthonormal basis of $(\ker F_*)^\perp$. Then $\{F_*(e_1), \dots, F_*(e_n)\}$ is an orthonormal basis of $\text{range} F_*$. By using (3.6) we can easily conclude that

$$\{F_*(e_1), \sec \theta \phi F_*(e_1), F_*(e_2), \sec \theta \phi F_*(e_2), \dots, F_*(e_p), \sec \theta \phi F_*(e_p)\}$$

is an orthonormal frame for $\Gamma(\text{range} F_*)$, where $2p = n = \text{rank} F_*$. Then we have the following result.

Lemma 3.1. *Let F be a slant Riemannian map from a Riemannian manifold (M_1, g_1) to an almost Hermitian manifold (M_2, g_2, J) with $\text{rank} F_* = n$. Let*

$$\{e_1, e_2, \dots, e_p\}, p = \frac{n}{2}$$

be a set of orthonormal vector fields in $(\ker F_*)^\perp$. Then

$$\{e_1, \sec \theta^* F_* \phi F_*(e_1), e_2, \sec \theta^* F_* \phi F_*(e_2), \dots, e_p, \sec \theta^* F_* \phi F_*(e_p)\}$$

is a local orthonormal basis of $(\ker F_*)^\perp$.

Proof. First by direct computation we have

$$g_1(e_i, \sec \theta^* F_* \phi F_*(e_i)) = \sec \theta g_2(F_*(e_i), \phi F_*(e_i)) = \sec \theta g_2(F_*(e_i), J F_*(e_i)) = 0.$$

In a similar way, we have

$$g_1(e_i, \sec \theta^* F_* \phi F_*(e_j)) = 0.$$

Since for a Riemannian map, we have ${}^*F_* \circ F_* = I$ (Identity map), by using (3.6), we get

$$g_1(\sec \theta^* F_* \phi F_*(e_i), \sec \theta^* F_* \phi F_*(e_j)) = \sec^2 \theta \cos^2 \theta g_1(e_i, e_j) = \delta_{ij},$$

which gives the assertion.

We now denote $*F_*\phi F_*$ by Q , then we have the following characterization of slant Riemannian maps.

Corollary 3.1. *Let F be a Riemannian map from a Riemannian manifold (M_1, g_1) to an almost Hermitian manifold (M_2, g_2, J) . Then F is a slant Riemannian map if and only if there exists a constant $\mu \in [-1, 0]$ such that*

$$Q^2X = \mu X$$

for $X \in \Gamma((\ker F_*)^\perp)$. If F is a slant Riemannian map, then $\mu = -\cos^2 \theta$.

Proof. By direct computation we have

$$Q^2X = *F_*\phi^2F_*(X).$$

Then proof comes from Theorem 3.1.

In the sequel we are going to show that the notion ω is useful to investigate the harmonicity of slant Riemannian map. To see this, we need the following Lemma.

Lemma 3.2. *Let F be a slant Riemannian map from a Riemannian manifold (M_1, g_1) to a Kähler manifold (M_2, g_2, J) . If ω is parallel then we have*

$$(\nabla F_*)(QX, QY) = -\cos^2 \theta (\nabla F_*)(X, Y) \quad (3.8)$$

for $X, Y \in \Gamma((\ker F_*)^\perp)$.

Proof. If ω is parallel, then from (3.4) we have

$$\mathcal{C}(\nabla F_*)(X, Y) = (\nabla F_*)(X, QY).$$

Interchanging the role of X and Y and taking into account that the second fundamental form is symmetric, we obtain

$$(\nabla F_*)(QX, Y) = (\nabla F_*)(X, QY).$$

Hence we get

$$(\nabla F_*)(QX, QY) = (\nabla F_*)(X, Q^2Y).$$

Then corollary 3.1 implies (3.8).

Theorem 3.2. *Let F be a slant Riemannian map from a Riemannian manifold $(M_1^{m_1}, g_1)$ to a Kähler manifold $(M_2^{m_2}, g_2, J)$. If ω is parallel then F is harmonic if and only if the distribution $(\ker F_*)$ is minimal.*

Proof. We choose an orthonormal basis of TM_1 as

$$\{v_1, \dots, v_r, e_1, \sec \theta Qe_1, e_2, \sec \theta Qe_2, \dots, e_s, \sec \theta Qe_s\}, \quad r + 2s = m_1$$

where $\{v_1, \dots, v_r\}$ is an orthonormal basis of $\ker F_*$ and $\{e_1, \sec \theta Qe_1, e_2, \sec \theta Qe_2, \dots, e_s, \sec \theta Qe_s\}$ is an orthonormal basis of $(\ker F_*)^\perp$. Since the second fundamental form is linear in every slot, we have

$$\tau = \sum_{i=1}^r (\nabla F_*)(v_i, v_i) + \sum_{j=1}^s (\nabla F_*)(e_j, e_j) + (\nabla F_*)(\sec \theta Qe_j, \sec \theta Qe_j).$$

Then from (3.8) we obtain

$$\tau = \sum_{i=1}^r (\nabla F_*)(v_i, v_i) = - \sum_{i=1}^r F_*(\nabla_{v_i} v_i)$$

which proves the assertion.

In the rest of this section, we investigate necessary and sufficient conditions for a slant Riemannian map to be totally geodesic. We recall that a differentiable map F between Riemannian manifolds (M_1, g_1) and (M_2, g_2) is called a totally geodesic map if $(\nabla F_*)(X, Y) = 0$ for all $X, Y \in \Gamma(TM_1)$.

Theorem 3.3. *Let F be a slant Riemannian map from a Riemannian manifold $(M_1^{m_1}, g_1)$ to a Kähler manifold $(M_2^{m_2}, g_2, J)$. Then F is totally geodesic if and only if*

- (i) *the fibres are totally geodesic,*
- (ii) *the horizontal distribution $(\ker F_*)^\perp$ is totally geodesic,*
- (iii) *For $X, Y \in \Gamma((\ker F_*)^\perp)$ and $V \in \Gamma((\text{range } F_*)^\perp)$ we have*

$$g_2(\mathcal{S}_{\omega F_*(Y)} F_*(X), \mathcal{B}V) = g_2(\nabla_X^{F^\perp} \omega F_*(Y), \mathcal{C}V) - g_2(\nabla_X^{F^\perp} \omega \phi F_*(Y), V).$$

Proof. For $X, Y \in \Gamma((\ker F_*)^\perp)$ and $V \in \Gamma((\text{range } F_*)^\perp)$, we have

$$g_2((\nabla F_*)(X, Y), V) = g_2(\nabla_X^F F_*(Y), V).$$

Then using (2.5), (3.1) and (3.2) we obtain

$$g_2((\nabla F_*)(X, Y), V) = -g_2(\nabla_X^F J\phi F_*(Y), V) + g_2(\nabla_X^F \omega F_*(Y), JV).$$

Using again (3.1), (3.2), (2.2) and Theorem 3.1, we get

$$\begin{aligned} g_2((\nabla F_*)(X, Y), V) &= \cos^2 \theta g_2(\nabla_X^F F_*(Y), V) - g_2(\nabla_X^{F^\perp} \omega \phi F_*(Y), V) \\ &\quad - g_2(\mathcal{S}_{\omega F_*(Y)} F_*(X), \mathcal{B}V) \\ &\quad + g_2(\nabla_X^{F^\perp} \omega F_*(Y), \mathcal{C}V) \end{aligned} \quad (3.9)$$

For $X, Y \in \Gamma((\ker F_*)^\perp)$ and $W \in \Gamma(\ker F_*)$, from (2.1) we derive

$$g_2((\nabla F_*)(X, W), F_*(Y)) = g_1(\nabla_X^1 Y, W). \quad (3.10)$$

In a similar way, for $U \in \Gamma(\ker F_*)$, we obtain

$$g_2((\nabla F_*)(U, W), F_*(Y)) = -g_1(\nabla_U^1 W, Y). \quad (3.11)$$

Then proof follows from (3.3), (3.9), (3.10) and (3.11).

4. Slant Riemannian maps and PHWC maps

In this section, we are going to show that every slant Riemannian map is a pseudo horizontally weakly conformal (PHWC) map, then we investigate the conditions for slant Riemannian maps to be pseudo horizontally homothetic map. First, we recall main definitions for PHWC maps. Let F be a map from a Riemannian manifold (M_1, g_1) to a Kähler manifold (M_2, g_2, J) , where g_1 and g_2 are Riemannian metrics on M_1 and M_2 , and J is the complex structure on M_2 . For any point $p \in M_1$, we denote the adjoint map of the tangent map $F_{*p} : T_p M_1 \longrightarrow T_{F(p)} M_2$ by ${}^*F_{*p} : T_{F(p)} M_2 \longrightarrow T_p M_1$. If $\text{range } F_{*p}$ is J -invariant, then we can define an almost complex structure $\tilde{J}_p$ on the horizontal space $(\ker F_{*p})^\perp$ by

$$\tilde{J}_p = F_{*p}^{-1} \circ J_{F(p)} \circ F_{*p}.$$

If the spaces $\text{range } F_{*p}$ are J -invariant for all p , then the almost complex structure on $(\ker F_*)^\perp$ is defined by

$$\tilde{J} = F_*^{-1} \circ J \circ F_*.$$

The map F is called pseudo-horizontally weakly conformal (PHWC) at p if and only if $\text{range } F_{*p}$ is J -invariant and $g_1|_{(\ker F_{*p})^\perp}$ is $\tilde{J}_p$ -Hermitian.

The map F is called pseudo-horizontally weakly conformal if and only if it is pseudo-horizontally weakly conformal at any point of p . A pseudo-horizontally weakly conformal map F from a Riemannian manifold (M_1, g_1) to a Kähler manifold (M_2, g_2, J) is called pseudo-horizontally homothetic if and only if $\tilde{J}$ is parallel in horizontal directions, i.e., $\nabla_X^1 \tilde{J} = 0$ for $X \in \Gamma((\ker F_*)^\perp)$ (for more details see [2], [3]).

Proposition 4.1. *Let F be a slant Riemannian map from a Riemannian manifold (M_1, g_1) to an almost Hermitian manifold (M_2, g_2, J) . Then F is PHWC map.*

Proof. We first note that $\tilde{J} = \sec \theta \phi$ is a complex structure on $(\text{range } F_*)$ and $\text{range } F_*$ is invariant with respect to $\tilde{J}$. Then we define $\hat{J} = \sec \theta Q = \sec \theta^* F_* \phi F_*$, it is easy to see that $\hat{J}$ is a complex structure on $(\ker F_*)^\perp$. Thus $((\ker F_*)^\perp, \hat{J})$ is an almost complex distribution. We now consider $\hat{g} = g_1|_{(\ker F_*)^\perp}$, then by direct computation we obtain

$$\hat{g}(\hat{J}X, \hat{J}Y) = \hat{g}(X, Y)$$

for $X, Y \in \Gamma((\ker F_*)^\perp)$. Thus $\hat{g}$ is $\hat{J}$ -Hermitian and $((\ker F_*)^\perp, \hat{g}, \hat{J})$ is almost Hermitian distribution. Therefore F is PHWC map.

From (3.4) and (2.1) we have the following.

Lemma 4.1. *Let F be a slant Riemannian map from a Riemannian manifold (M_1, g_1) to a Kähler manifold (M_2, g_2, J) . Then ϕ is parallel if and only if*

$$(\nabla F_*)(X, QY) = \nabla_X^F \phi F_*(Y) - \phi F_*(\nabla_X^1 Y)$$

for $X, Y \in \Gamma((\ker F_*)^\perp)$.

We now give necessary and sufficient conditions for a slant Riemannian map to be pseudo horizontally homothetic map.

Theorem 4.1. *Let F be a slant Riemannian map from a Riemannian manifold (M_1, g_1) to a Kähler manifold (M_2, g_2, J) . Then F is pseudo horizontally homothetic map if and only if ϕ is parallel and*

$$(\nabla F_*)(X, U) = 0$$

for $X \in \Gamma((\ker F_*)^\perp)$ and $U \in \Gamma(\ker F_*)$.

Proof. By direct computations we have

$$(\nabla_X \hat{J})Y = \sec \theta \nabla_X^1 QY - Q \nabla_X^1 Y.$$

Hence we obtain

$$F_*(\nabla_X \hat{J})Y = \sec \theta (F_*(\nabla_X^1 QY) - \phi F_*(\nabla_X^1 Y)).$$

Then using (2.1) we get

$$F_*(\nabla_X \hat{J})Y = \sec \theta (-(\nabla F_*)(X, QY) + \nabla_X^F \phi F_*(Y) - \phi F_*(\nabla_X^1 Y)). \quad (4.1)$$

On the other hand, since QY and $U \in \Gamma(Ker F_*)$ are orthogonal, we have

$$g_1((\nabla_X^1 \hat{J})Y, U) = \sec \theta g_1(QY, \nabla_X^1 U).$$

Then adjoint map $*F_*$ and (2.1) imply

$$g_1((\nabla_X^1 \hat{J})Y, U) = \sec \theta g_2(\phi F_*(Y), (\nabla F_*)(X, U)). \quad (4.2)$$

Then proof comes from (4.1) and (4.2).

Remark 4.1. A harmonic map which is pseudo horizontally weakly is called a pseudo harmonic morphism. Pseudo harmonic morphisms were defined in [15] by Loubeau. He also obtained that a smooth map F from a Riemannian manifold M to a Kähler manifold N is a pseudo-harmonic morphism if and only if F pulls back local pluriharmonic functions on N to local harmonic functions on M . In this paper, we show that any slant Riemannian map is a pseudo horizontally weakly conformal map (Proposition 4.1). Also Theorem 3.2 of the present paper shows that it is possible to obtain harmonic slant Riemannian maps under some geometric conditions. Thus slant Riemannian maps are good candidates for pseudo harmonic morphisms.

Acknowledgment. This paper is supported by The Scientific and Technological Council of Turkey (TUBITAK) with number (TBAG-109T125).

References

- [1] Abraham, R., Marsden, J.E., Ratiu, T., *Manifolds, Tensor Analysis, and Applications*, Springer-Verlag, Newyork, 1988.
- [2] Aprodu, M.A., Aprodu, A. M., Implicitly defined harmonic PHH submersions, *Manuscripta Math.* 100,(1999), 103-121.

- [3] Aprodu, M.A., Aprodu, A. M., Brînzănescu, A class of harmonic maps and minimal submanifolds, *Int. J. Math.* 11 (2000), 1177-1191.
- [4] Baird, P. and Wood, J. C. *Harmonic Morphisms Between Riemannian Manifolds*, London Mathematical Society Monographs, No. 29, Oxford University Press, The Clarendon Press, Oxford, 2003.
- [5] Burns, D., Burstall, F. E., de Bartolomeis, P., Rawnsley, J., Stability of harmonic maps of Kähler manifolds. *J. Diff. Geom.* 30, (1989), 579-594.
- [6] Cabrerizo, J. L., Carriazo, A., Fernandez, L. M., Fernandez, M., Slant submanifolds in Sasakian manifolds. *Glasg. Math. J.* 42(1), (2000), 125-138.
- [7] Chen, B. Y., Slant immersions. *Bull. Austral. Math. Soc.* 41 (1), (1990), 135-147.
- [8] Chen, B. Y., *Geometry of slant Submanifolds*, Katholieke Universiteit Leuven, Leuven, 1990.
- [9] Chinea, D., Almost contact metric submersions. *Rend. Circ. Mat. Palermo*, (1985), 34(1), 89-104.
- [10] Falcitelli, M., Ianus, S., Pastore, A. M., *Riemannian Submersions and Related Topics*. World Scientific, River Edge, NJ, 2004.
- [11] Fischer, A. E.: Riemannian maps between Riemannian manifolds, *Contemporary math.* 132, 331-366, (1992).
- [12] Garcia-Rio, E., Kupeli, D. N., *Semi-Riemannian maps and their Applications*, Kluwer Academic, Dordrecht, 1999.
- [13] Gray, A., Pseudo-Riemannian almost product manifolds and submersions, *J. Math. Mech.*, (1967), 16, 715-737.
- [14] Ianus, S., Mazzocco, R., Vilcu, G. E., Riemannian submersions from quaternionic manifolds. *Acta Appl. Math.*, (2008), 104(1), 83-89.
- [15] Loubeau, E. Pseudo-harmonic morphisms, *Internat. J. Math.* 8, no. 7, (1997), 943-957.
- [16] Loubeau, E., Slobodeanu, R., Eigenvalues of harmonic almost submersions, *Geom Dedicata* 145, (2010), 103-126

- [17] Marrero, J. C., Rocha, J., Locally conformal Kähler submersions. *Geom. Dedicata*, (1994), 52(3), 271-289.
- [18] Nore, T., Second fundamental form of a map, *Ann. Mat. Pur. and Appl.*, 146, (1987), 281-310.
- [19] O'Neill, B., The fundamental equations of a submersion, *Mich. Math. J.*, (1966), 13, 458-469.
- [20] Şahin, B., Invariant and anti-invariant Riemannian maps to Kähler manifolds, *Int. J. Geom. Methods Mod. Phys.*, vol:7, no:3 (2010), 1-19.
- [21] Şahin, B., Semi-invariant Riemannian maps to Kähler manifolds, *Int. J. Geom. Methods Mod. Phys.*, 8(7), (2011), 1439-1454.
- [22] Watson, B. Almost Hermitian submersions. *J. Differential Geometry*, (1976), 11(1), 147-165.
- [23] Yano, K. and Kon, M. *Structures on Manifolds*, World Scientific, Singapore, 1984.